

Pacira Reports Preliminary Net Product Sales of \$118.7 Million for the First Quarter of 2021

April 7, 2021

EXPAREL average daily sales at 115% of the prior year for the first quarter and 137% of the prior year for the month of March

PARSIPPANY, N.J., April 07, 2021 (GLOBE NEWSWIRE) -- Pacira BioSciences, Inc. (Nasdaq: PCRX), the industry leader in its commitment to non-opioid pain management and regenerative health solutions, today reported preliminary unaudited net product sales of EXPAREL[®] (bupivacaine liposome injectable suspension) and iovera° of \$114.7 million and \$3.3 million, respectively, for the first quarter of 2021. The company also reported preliminary unaudited net product sales of EXPAREL and iovera° of \$44.3 million and \$1.5 million, respectively, for the month of March 2021. EXPAREL average daily sales were 108%, 103%, and 137% of the prior year levels for the months of January, February and March, respectively.

"The first quarter of 2021 was marked by significant progress as highlighted by the U.S. regulatory approval of EXPAREL in the pediatric setting and its growing utilization across a wide range of long-acting regional blocks for low or no opioid postsurgical pain control," said Dave Stack, chairman and chief executive officer of Pacira BioSciences. "The surgical paradigm continues its rapid progression out of the hospital as EXPAREL-based enhanced recovery after surgery protocols redefine best practice and drive surgical migration to the outpatient setting."

"The COVID-19 pandemic has accelerated this transition, which we expect will be permanently embedded in healthcare systems given the improved patient outcomes, satisfaction rates and economic advantages of outpatient care. These positive market dynamics, along with EXPAREL growth continuing to outpace the recovery of the elective surgery market, leave us very well-positioned for robust long-term growth once the COVID-19 crisis recedes and elective surgeries normalize," added Mr. Stack.

The company's 2021 product sales continue to be negatively impacted by the COVID-19 pandemic, which mandated significant postponement or suspension in the scheduling of elective surgical procedures resulting from public health guidance and government directives. Elective surgery restrictions began to lift on a state-by-state basis in April 2020. In order to provide greater transparency, the company will continue to report monthly intra-quarter unaudited net product sales until it has gained enough visibility around the impacts of COVID-19. The company is also providing weekly EXPAREL utilization and elective surgery data within its investor presentation, which is accessible at investor pacina.com.

The financial information included in this press release is preliminary, unaudited and subject to adjustment. It does not present all information necessary for an understanding of the company's financial results for the first quarter or full year 2021.

About Pacira BioSciences

Pacira BioSciences, Inc. (Nasdaq: PCRX) is the industry leader in its commitment to non-opioid pain management and regenerative health solutions to improve patients' journeys along the neural pain pathway. The company's long-acting local analgesic, EXPAREL [®] (bupivacaine liposome injectable suspension) was commercially launched in the United States in April 2012. EXPAREL utilizes DepoFoam[®], a unique and proprietary product delivery technology that encapsulates drugs without altering their molecular structure, and releases them over a desired period of time. In April 2019, Pacira acquired the iovera° system, a handheld cryoanalgesia device used to deliver precise, controlled doses of cold temperature only to targeted nerves. To learn more about Pacira, including the corporate mission to reduce overreliance on opioids, visit www.pacira.com.

About EXPAREL®

EXPAREL (bupivacaine liposome injectable suspension) is indicated in patients 6 years of age and older for single-dose infiltration to produce postsurgical local analgesia, and in adults as an interscalene brachial plexus nerve block to produce postsurgical regional analgesia. Safety and efficacy have not been established in other nerve blocks. The product combines bupivacaine with DepoFoam®, a proven product delivery technology that delivers medication over a desired time period. EXPAREL represents the first and only multivesicular liposome local anesthetic that can be utilized in the peri- or postsurgical setting. By utilizing the DepoFoam platform, a single dose of EXPAREL delivers bupivacaine over time, providing significant reductions in cumulative pain scores with up to a 78 percent decrease in opioid consumption; the clinical benefit of the opioid reduction was not demonstrated. Additional information is available at www.EXPAREL.com.

Important Safety Information for Patients

EXPAREL should not be used in obstetrical paracervical block anesthesia. In studies in adults where EXPAREL was injected into a wound, the most common side effects were nausea, constipation, and vomiting. In studies in adults where EXPAREL was injected near a nerve, the most common side effects were nausea, fever, and constipation. In the study where EXPAREL was given to children, the most common side effects were nausea, vomiting, constipation, low blood pressure, low number of red blood cells, muscle twitching, blurred vision, itching, and rapid heartbeat. EXPAREL can cause a temporary loss of feeling and/or loss of muscle movement. How much and how long the loss of feeling and/or muscle movement depends on where and how much of EXPAREL was injected and may last for up to 5 days. EXPAREL is not recommended to be used in patients younger than 6 years old for injection into the wound, for patients younger than 18 years old for injection near a nerve, and/or in pregnant women. Tell your health care provider if you or your child has liver disease, since this may affect how the active ingredient (bupivacaine) in EXPAREL is eliminated from the body. EXPAREL should not be injected into the spine, joints, or veins. The active ingredient in EXPAREL can affect the nervous system and the cardiovascular system; may cause an allergic reaction; may cause damage if injected into the joints; and can cause a rare blood disorder.

About iovera®

The iovera° system is used to destroy tissue during surgical procedures by applying freezing cold. It can also be used to produce lesions in peripheral nervous tissue by the application of cold to the selected site for the blocking of pain. It is also indicated for the relief of pain and symptoms associated with osteoarthritis of the knee for up to 90 days. In one study, the majority of the patients suffering from osteoarthritis of the knee experienced pain and

system relief beyond 150 days.¹ The iovera° system's "1×90" Smart Tip configuration (indicating one needle which is 90 mm long) can also facilitate target nerve location by conducting electrical nerve stimulation from a separate nerve stimulator. The iovera° system is not indicated for treatment of central nervous system tissue.

Important Safety Information

The iovera° system is contraindicated for use in patients with the following: Cryoglobulinemia; Paroxysmal cold hemoglobinuria; cold urticaria; Raynaud's disease; open and/or infected wounds at or near the treatment line. Potential complications: As with any surgical treatment that uses needle-based therapy, there is potential for temporary site-specific reactions, including but not limited to: bruising (ecchymosis); swelling (edema); inflammation and/or redness (erythema); pain and/or tenderness; altered sensation (localized dysesthesia). Typically, these reactions resolve with no physician intervention. Patients may help the healing process by applying ice packs to the affected sites, and by taking over-the-counter analgesics.

Forward-Looking Statements

Any statements in this press release about the company's future expectations, plans, trends, outlook, projections and prospects, and other statements containing the words "believes," "anticipates," "estimates," "expects," "intends," "may," "will," "would," "could," "can" and similar expressions, constitute forward-looking statements within the meaning of The Private Securities Litigation Reform Act of 1995. Actual results may differ materially from those indicated by such forward-looking statements as a result of various important factors, including risks relating to: the impact of the worldwide COVID-19 (Coronavirus) pandemic and related global economic conditions; the success of the company's sales and manufacturing efforts in support of the commercialization of EXPAREL; the rate and degree of market acceptance of EXPAREL; the size and growth of the potential markets for EXPAREL and the company's ability to serve those markets; the company's plans to expand the use of EXPAREL to additional indications and opportunities, and the timing and success of any related clinical trials; the ability to realize anticipated benefits and synergies from the acquisition of MyoScience; the ability to successfully integrate iovera° and any other future acquisitions into the company's existing business; the commercial success of iovera° and other factors discussed in the "Risk Factors" of the company's most recent Annual Report on Form 10-K and in other filings that the company periodically makes with the SEC. In addition, the forward-looking statements included in this press release represent the company's views as of the date of this press release. Important factors could cause actual results to differ materially from those indicated or implied by forward-looking statements, and as such the company anticipates that subsequent events and developments will cause its views to change. However, while the company may elect to update these forward-looking statements at some point in the future, it specifically disclaims any obligat

Investor Contact: Susan Mesco, (973) 451-4030 susan.mesco@pacira.com Media Contact: Coyne Public Relations Alyssa Schneider, (973) 588-2270 aschneider@coynepr.com

¹ Radnovich, R. et al. "Cryoneurolysis to treat the pain and symptoms of knee osteoarthritis: a multicenter, randomized, double-blind, sham-controlled trial." Osteoarthritis and Cartilage (2017) p1-10.